
NAME _____

- 1. A 51.2 g of O_2 gas (molecular weight 32 g/mole) occupies _____ L at 22 °C and 2.5 atm. Use the ideal gas law, PV = nRT
 - P: pressure (atm)
 - V: volume (L)
 - n: # of moles
 - R: $0.082 (atm \cdot L)/(mole \cdot K)$
 - T: temperature (K)
 - a. 0.079
- b. 3.2 x 10⁻²
- c. 15.5
- d. 1.2
- e. 12.6
- 2. Arrange the following gases in order of increasing average molecular speed at 25 °C.
 - He, O₂, CO₂, H₂
 - a. $He < O_2 < H_2 < CO_2$
- b. $CO_2 < O_2 < He < H_2$
- c. $He < H_2 < O_2 < CO_2$
- d. $CO_2 < He < N_2 < O_2$
- e. $CO_2 < O_2 < H_2 < He$
- 3. Which one of the following is the correct electron configuration for a ground-state carbon atom?



- 4. The lines in the absorption spectrum of hydrogen result from ______.
 - a. electrons given off by hydrogen as it cools

b. energy given off in the form of visible light when an electron moves from a higher energy

	c. electrons gived. energy takento a higher er	in the form of visi		uminated by visible lig etron moves from a low					
5.	The wavelength <i>c</i> (speed of light)	m.							
	a. 0.0400 b.	12.0	c. 2.5	d. 2.50×10^{-5}	e. 25.0				
6.	The quantum number which describes the shape of an orbital is:								
	a. ne. The shape of	b. <i>l</i> an atomic orbital	c. m_l is nothing to do with	d. <i>ms</i> quantum number					
7.	There are possible values for the magnetic quantum number (m_l) of an electron in a 3p subshell.								
	a. 0	b. 1	c. 2	d. 3	e. 4				
8.	For a d orbital,								
	b. The value ofc. The value of	m_l must be 3							
9.	not possible for a	an electron in an at m_s 1 m_s 1 $\frac{1}{2}$ 1 $\frac{1}{2}$ 2 $\frac{1}{2}$ 2 $-\frac{1}{2}$, and m_s , which one of	these sets is				
10.	Which atom belo	ow is diamagnetic?	?						
	a. Be	b. B	c. C	d. N	e. O				
Consi	(i) [Kr] 5s ¹ (ii) [Ne] 3s ² 3p ⁵ (iii) [Ar] 4s ² 3d ¹⁰ (iv) [Ne] 3s ² 3p ⁶) 4p ⁴	tions to answer the que	estions 11 and 12 that	follow:				

(v) [Ar] $4s^2$

11. The electron configuration of the atom that is expected to form a stable +2 ion is ______.

a. (i)

b. (ii)

c. (iii)

d. (iv)

e. (v)

12. The electron configurations of the two atoms that form isoelectronic ions are ______.

a. (i) and (iii)

b. (ii) and (iii)

c. (ii) and (v)

d. (iii) and (iv)

e. (i) and (v)

13. The ion PCl₄ has ______ valence electrons.

a. 34

b. 8

c. 28

d. 35

e. 36

14. Of the following, which gives the correct order for atomic radius for Mg, Na, P, Si and Ar?

a. Si > P > Ar > Na > Mg

b. Na > Mg > Si > P > Ar

c. Ar > P > Si > Mg > Na

d. Ar > Si > P > Na > Mg

e. Mg > Na > P > Si > Ar

15. Which isoelectronic series is correctly arranged in order of increasing radius?

a. $Ca^{2+} < K^+ < Ar < Cl^-$


b. $Cl^{-} < Ar < K^{+} < Ca^{2+}$

 $c. \quad Ca^{2+} < Ar < K^+ < Cl^-$

d. $Ca^{2+} < K^+ < Cl^- < Ar$

e. $K^+ < Ca^{2+} < Ar < Cl^{-1}$

16. The formal charge on **sulfur** of SO₄²⁻ ion in figure is _____



a. +4

b. -2

c. +2

d. +0

e. -4

17. Given the electronegativities below, which covalent single bond is least polar?

Element:

Electronegativity:

H 2.1 C 2.5 N 3.0 O 3.5

b. O-C

b. O-N

c. N-H

d. C-H

e. O-H

18. According to valence bond theory, which orbitals overlap in the formation of the bond in F_2 ?

- a. 1s on F and 1s on F
- b. 1s on F and 2s on F
- c. 2s on F and 2s on F
- d. 2s on F and 2p on F
- e. 2p on F and 2p on F
- 19. The basis of the valence shell electron pair repulsion (VSEPR) model of molecular bonding is
 - a. regions of electron density in the valence shell of an atom will arrange themselves so as to maximize overlap
 - b. atomic orbitals of the bonding atoms must overlap for a bond to form
 - c. electron domains in the valence shell of an atom will arrange themselves so as to minimize repulsions
 - d. hybrid orbitals will form as necessary to, as closely as possible, achieve spherical symmetry
 - e. regions of electron density on an atom will organize themselves so as to maximize scharacter
- 20. What is the orbital hybridization of the central atom P in PF₅?
 - a. sp
 - b. sp^2
 - c. sp^3
 - d. sp^3d
 - e. sp^3d^2
- 21. An energy level scheme for the orbitals of second row diatomic molecules O₂ through Ne₂, lists the molecular orbitals in the following order of increasing energy

$$\sigma_{1s} \, < \, \sigma *_{1s} \, < \sigma_{2s} \, < \, \sigma *_{2s} \, < \, \sigma_{2p(z)} \, < \, \pi_{2p(y)}, \, \pi_{2p(x)} \, < \, \pi *_{2p(y)}, \, \pi *_{2p(x)} \, < \, \sigma *_{2p(z)}$$

Based on this energy level scheme, the bond order for the bond in the O_2 ion in its ground state is

- a. 0.5
- b. 1.0
- c. 1.5
- d. 2.0
- e. 2.5
- 22. Find partial pressures of He (P_{He}) and O_2 (P_{O2}) in a 12.5 L tank with 12.5 g He and 8.6 g O_2 at 298 K (10 points). Ideal gas law, PV = nRT

P: pressure (atm)

V: volume (L)

N: # of moles

R: $0.082 (atm \cdot L)/(mole \cdot K)$

	: temperature (K)
23.	For Cu (atomic number 29), (15 pts)
	a. Write the electron configuration of Cu (atomic number 29) (5 pts) (energy level of atomic orbitals: 1s, 2s, 2p, 3s, 3p, 4s, 3d, 4p)
	b. How many valence electrons? (5 pts)
	c. Diamagnetic or paramagnetic? (5 pts)
	Draw the Lewis dot structures of the following compounds in the column (1), using VSEF model, predict the molecular geometry of the compound in the column (2), and decide whether the molecule is polar or nonpolar. You can use a line to represent a pair of bonding electrons between atoms. Missing lone-pair electrons of peripheral atoms in the Lewis d

diagram will result in a deduction of 2 points each occurrence. (45 pts)

Put the keys in the box directly

	Lewis structure (5 pts)	Molecular geometry (5 pts)	Polar or Nonpolar (5 pts)
CCl ₄	: Cl : : Cl : C : Cl : : Cl :		
OCl ₂	:CI • O • CI:		
XeF ₄	Xe F		